

Forward Compton Scattering, using Real Analytic Amplitudes

Martin M. Block *

*Department of Physics and Astronomy,
Northwestern University, Evanston, IL 60208*

Abstract

We analyze forward Compton scattering, using real analytic amplitudes. By fitting the total γp scattering cross section data in the high energy region $5 \text{ GeV} \leq \sqrt{s} \leq 200 \text{ GeV}$, using a cross section rising as $\ln^2 s$, we calculate $\rho_{\gamma p}$, the ratio of the real to the imaginary portion of the forward Compton scattering amplitude, and compare this to ρ_{nn} , the ratio of the *even* portions of the pp and $\bar{p}p$ forward scattering amplitudes. We find that the two ρ -values are, within errors, the same in the c.m.s. energy region $5 \text{ GeV} \leq \sqrt{s} \leq 200 \text{ GeV}$, as predicted by a factorization theorem of Block and Kadailov[5].

*Work partially supported by Department of Energy contract DA-AC02-76-Er02289 Task D.

1 Introduction

The ρ -value is defined as the ratio of the real to the imaginary portion of the forward scattering amplitude. Block and Kadailov[5] have shown that $\rho_{nn} = \rho_{\gamma p}$ if one uses eikonals for the even portion of nucleon-nucleon scattering and γp scattering that have *equal* opacities, *i.e.*, eikonals that have the same value at impact parameter $b = 0$. This is the equivalent of the more physical statement that

$$\left(\frac{\sigma_{\text{elastic}}(s)}{\sigma_{\text{tot}}(s)} \right)_{\gamma p} = \left(\frac{\sigma_{\text{elastic}}(s)}{\sigma_{\text{tot}}(s)} \right)_{nn}, \quad \text{for all } s. \quad (1)$$

Block and Kaidalov[5] have proved three factorization theorems:

$$1. \quad \frac{\sigma_{nn}(s)}{\sigma_{\gamma p}(s)} = \frac{\sigma_{\gamma p}(s)}{\sigma_{\gamma\gamma}(s)},$$

where the σ 's are the total cross sections for nucleon-nucleon, γp and $\gamma\gamma$ scattering,

$$2. \quad \frac{B_{nn}(s)}{B_{\gamma p}(s)} = \frac{B_{\gamma p}(s)}{B_{\gamma\gamma}(s)},$$

where the B 's are the nuclear slope parameters for elastic scattering,

$$3. \quad \rho_{nn}(s) = \rho_{\gamma p}(s) = \rho_{\gamma\gamma}(s),$$

where the ρ 's are the ratio of the real to imaginary portions of the forward scattering amplitudes,

with the first two factorization theorems having their own proportionality constant. These theorems are exact, for *all* s (where $\sqrt{s}$ is the c.m.s. energy), and survive exponentiation of the eikonal[5]. The last theorem is valid independently of the model which takes one from nn to γp to $\gamma\gamma$ reactions, as long as the respective eikonals have equal opacities, *i.e.*, eq. (1) holds. We wish to demonstrate experimentally here the validity of the theorem that states that $\rho_{nn}(s) = \rho_{\gamma p}(s)$. However, no data are available in the hadronic sector for $\rho_{\gamma p}$. The purpose of this note is to analyze forward Compton scattering at high energies in order to extract $\rho_{\gamma p}$ and then to compare it to ρ_{nn} .

Damashek and Gilman[3] in 1970 have calculated $\rho_{\gamma p}$ using a singly-subtracted dispersion relation, up to a gamma ray laboratory energy $\nu = 20$ GeV, *i.e.*, to a c.m.s. energy of $\sqrt{s} = 6.2$ GeV. Here we extend the $\rho_{\gamma p}$ evaluation up to $\sqrt{s} = 200$ GeV and then compare the results to ρ_{nn} , the ratio of the *even* portions of the $\bar{p}p$ and pp forward scattering amplitudes (for references, see [4]). In this paper we will calculate $\rho_{\gamma p}$ using real analytic amplitudes (see Section G. p.583 of ref. [1]). It is shown in ref. [1] that the numerical complexities of dispersion relations in analyzing $\bar{p}p$ and pp scattering can be circumvented by direct use of analytic functions to fit forward $\bar{p}p$ and pp scattering amplitudes, a technique first proposed by Bourrely and Fischer[2]. We will introduce a variant of the Block and Cahn analysis[1] appropriate for Compton scattering, $\gamma p \rightarrow \gamma p$.

2 Preliminaries

This work largely follows the procedures and conventions used by Block and Cahn[1]. We use units where $\hbar = c = 1$. The variable s is the square of the c.m. system energy, whereas ν is the laboratory system momentum. In terms of the *even* laboratory scattering amplitude f_+ , where $f_+(\nu) = f_+(-\nu)$, the total unpolarized Compton cross section σ_{tot} is given by[3]

$$\sigma_{\text{tot}} = \frac{4\pi}{\nu} \text{Im} f_+(\theta = 0), \quad (2)$$

where θ is the laboratory scattering angle. We will assume that our amplitudes are real analytic functions with a simple cut structure[1]. For γp scattering, the assumed cut structure is a left-hand cut that begins at the gamma ray energy $-\nu_0$ and a right-hand cut that begins at the gamma ray energy ν_0 , with a real amplitude on the real axis between $-\nu_0$ and ν_0 . The threshold gamma ray energy for pion production is $\nu_0 = m_\pi + \frac{m_\pi^2}{m} \approx 0.16$ GeV, where m_π and m are the pion and proton masses, respectively. We will use an even amplitude for γp reactions in the high energy region $\nu \gg \nu_0$, far above any cuts, (see ref.[1], p. 587, eq. (5.5a), with $a = 0$), where the even amplitude simplifies considerably and is given by

$$f_+ = i \frac{\nu}{4\pi} \left\{ A + \beta [\ln(s/s_0) - i\pi/2]^2 + c s^{\mu-1} e^{i\pi(1-\mu)/2} \right\} + C_{\text{subtraction}}, \quad (3)$$

where A , β , c , s_0 and μ are real constants. The additional real constant $C_{\text{subtraction}} = f_+(0)$ is the subtraction constant at $\nu = 0$ needed in a singly-subtracted dispersion relation[3] for the reaction $\gamma + p \rightarrow \gamma + p$ and is given by the Thompson scattering limit, *i.e.*, $f_+(0) = -\alpha/m = -3.03 \mu\text{b GeV}$. In eq. (3), we have assumed that the Compton cross section rises as $\ln^2(s)$ at ultra-high energies.

The real and imaginary parts of eq. (3) are given by

$$\text{Re} \frac{4\pi}{\nu} f_+ = \beta \pi \ln s/s_0 - c \cos(\pi\mu/2) s^{\mu-1} + \frac{4\pi}{\nu} f_+(0) \quad (4)$$

$$\text{Im} \frac{4\pi}{\nu} f_+ = A + \beta \left[\ln^2 s/s_0 - \frac{\pi^2}{4} \right] + c \sin(\pi\mu/2) s^{\mu-1}, \quad (5)$$

where $s = 2m\nu + m^2 \approx 2m\nu$ and with c being a real constant. Using equations (2), (4) and (5), we find the total cross section for high energy Compton scattering is given by

$$\sigma_{\text{tot}} = A + \beta \left[\ln^2 s/s_0 - \frac{\pi^2}{4} \right] + c \sin(\pi\mu/2) s^{\mu-1}, \quad (6)$$

and that ρ , the ratio of the real to the imaginary portion of the forward scattering amplitude, is given by

$$\rho = \frac{\beta \pi \ln s/s_0 - c \cos(\pi\mu/2) s^{\mu-1} + \frac{4\pi}{\nu} f_+(0)}{\sigma_{\text{tot}}}, \quad (7)$$

with $f_+(0) = -3.03 \mu\text{b GeV}$. We will use units of ν in GeV and s in GeV^2 , and cross sections in μb . We have to fit the 5 real constants A , c , β , s_0 and μ . If we assume that the term in c is a Regge descending term, then $\mu = 1/2$.

The high energy behavior that was assumed by Damashek and Gilman[3] in 1970 when they calculated $\rho_{\gamma p}$ numerically using dispersion relations was that the cross section approached a constant value asymptotically, *i.e.*, they assumed that σ_{tot} , for $\nu \rightarrow \infty$, was given by

$$\sigma_{\text{tot}} = A + \frac{c'}{\nu^{1/2}}, \quad (8)$$

with $A = 96.6\mu\text{b}$ and $c' = 70.2\mu\text{b GeV}^{1/2}$, with ν measured in GeV. However, today we know experimentally that the cross section rises at high energies and that the rising term can be fit by a $\ln^2 s$ term, as expressed in eq. (6).

As a check on our real analytic amplitude analysis, the highest energy ρ values of ref. [3], where Damashek and Gilman used dispersion relations assuming the asymptotic cross section of eq. (8), can be simply reproduced from eq. (7) with $\beta = 0$ and eq. (8) by the relation

$$\rho_{\text{analytic}} = -\frac{70.2/\nu^{1/2} + 38.07/\nu}{96.6 + 70.2/\nu^{1/2}}. \quad (9)$$

The ρ -values calculated from eq. (9) and the results from the singly-subtracted dispersion relation of ref. [3] agree to better than 2% over the energy range $10 \text{ GeV} \leq \nu \leq 20 \text{ GeV}$. The numerical agreement is excellent—the simplicity and ease of calculation at high energies using real analytical amplitudes compared to a dispersion relation analysis is clear.

The ρ values in ref. [3] were only calculated up to $\nu = 20 \text{ GeV}$, which corresponds to a c.m.s. energy of $\sqrt{s} = 6.2 \text{ GeV}$. We now make an amplitude analysis for $\rho_{\gamma p}$ using a *rising* cross section, asymptotically going as $\ln^2 s$, by fitting the experimental cross sections σ_{tot} in the energy interval $5 \text{ GeV} \leq \sqrt{s} \leq 200 \text{ GeV}$ to the parameters A , c , β and s_0 of eq. (6), using a Regge descending trajectory with $\mu = 0.5$.

3 Results and Conclusions

Since cross sections for γp scattering are now available for c.m.s. energies up to 200 GeV, we made a χ^2 fit to the experimental $\sigma_{\text{tot}}(\gamma p)$ data in the c.m.s. energy interval $5 \text{ GeV} \leq \sqrt{s} \leq 200 \text{ GeV}$. We find a reasonable representation of the data using eq. (6), with a χ^2 per degree of freedom of 0.98 for 40 degrees of freedom, with the coefficients:

$A = 115 \pm 13$, $c = 35.1 \pm 97$, $\beta = 1.07 \pm 0.57$, $s_0 = 68.4 \pm 80.2 \text{ GeV}^2$, using a fixed value of $\mu = 0.5$ (cross sections in μb when s is in GeV^2). This fit, plotted as a function of c.m.s. energy, gives the dashed cross section curve $\sigma_{\text{tot}}(\gamma p)$ in Fig. 1, as well as the dashed $\rho_{\gamma p}$ curve in Fig. 2, using eq. (6) and eq. (7), respectively. Several remarks are in order. The experimental data are taken from the Particle Data Group compilation[6] and include the only three high energy points—in the neighborhood of 200 GeV—that are available. The very large errors of β and s_0 clearly reflect the fact that these few points also have large systematic errors ranging from ≈ 7 to 21%. The accuracy is sufficient to show that the cross section rises, but not much more. Obviously, much more precise data at high energies are required for pinning down the $\log^2 s$ parameters.

In order to show visually the sensitivity of $\sigma_{\text{tot}}(\gamma p)$ and $\rho_{\gamma p}$ to the parameters of the fit, we have also plotted in Fig. 1 the dotted curve (a slight variation of the parameters of A and c within their errors), where we have set $A = 107$ and $c = 134$. This curve has as its $\rho_{\gamma p}$ analog the dotted curve of Fig. 2. Using eq. (7), Fig. 2 shows our result for $\rho_{\gamma p}$ compared to ρ_{nn} , the ρ -value for nucleon-nucleon scattering found in ref. [4], as a function of the c.m.s. energy $\sqrt{s}$, in GeV. The solid curve is ρ_{nn} ; the dashed line is the $\rho_{\gamma p}$ curve which corresponds to the central values $A = 115$, $c = 35.1$, $\beta = 1.07$, $s_0 = 68.4 \text{ GeV}^2$; the dotted line is the $\rho_{\gamma p}$ curve which uses the slightly varied parameters $A = 107$ and $c = 134$. The agreement between the slightly modified $\rho_{\gamma p}$ and ρ_{nn} over the energy interval $5 \text{ GeV} \leq \sqrt{s} \leq 200 \text{ GeV}$ lends experimental support, in a model independent way, for the three factorization theorems of Block and Kadailov[5, 7]. Of course, precision cross section data in the region $20 \text{ GeV} < \sqrt{s} < 200 \text{ GeV}$ would enable us to strengthen this conclusion.

4 Acknowledgments

The author would like to thank Professor J. D. Jackson for valuable critical comments and suggestions.

References

- [1] M. M. Block and R. N. Cahn, Rev. Mod. Phys. **57**, 563 (1985).
- [2] C. Bourrely and J. Fischer, Nucl. Phys. B **61**, 513 (1973).
- [3] M. Damashek and F. Gilman, Phys. Rev. D **1**, 1319 (1970).
- [4] M. M. Block *et al.*, e-Print Archive: **hep-ph/0004232**, Phys. Rev. D **62**, 077501 (2000).
- [5] M. M. Block and A. B. Kaidalov, e-Print Archive: **hep-ph/0012365**, Phys. Rev. D **64**, 076002 (2001).
- [6] D. E. Groom *et al.*, Eur. Phys. J. C **15**, 1 (2000).
- [7] M. M. Block *et al.*, e-Print Archive: **hep-ph/0111046 v2**, Eur. Phys. J. C **23**, 329 (2002).




Figure 1: The dashed curve is $\sigma_{\text{tot}}(\gamma p)$, the predicted total γp cross section from eq. (6), using the central value parameters $A = 115$, $c = 35.1$, $\beta = 1.07$, $s_0 = 68.4 \text{ GeV}^2$, and $\mu = 0.5$ of a χ^2 fit, compared to the existing high energy experimental data in the c.m.s. energy interval $5 \text{ GeV} \leq \sqrt{s} \leq 200 \text{ GeV}$. The dotted curve varies the parameters slightly, with $A \rightarrow 107$ and $c \rightarrow 134$ (values within their errors). The corresponding $\rho_{\gamma p}$ curves are shown in Fig. 2.




Figure 2: The solid curve is ρ_{nn} , the predicted ratio of the real to imaginary part of the forward scattering amplitude for the ‘elastic reactions’, $\gamma + p \rightarrow V + p$ scattering amplitude, where V is ρ , ω or ϕ (using factorization). The dashed curve is $\rho_{\gamma p}$, the ratio of the real to imaginary part of the forward scattering amplitude for Compton scattering, $\gamma + p \rightarrow \gamma + p$, found from eq. (7), using real analytic amplitudes that asymptotically go as $\ln^2 s$, with best fit parameters $A = 115$, $c = 35.1$, $\beta = 1.07$, $s_0 = 68.4 \text{ GeV}^2$, and $\mu = 0.5$. The dotted line is the $\rho_{\gamma p}$ curve where the parameters of the fit have been slightly varied within their errors, with $A \rightarrow 107$ and $c \rightarrow 134$. The corresponding two curves for $\sigma_{\text{tot}}(\gamma p)$ are shown in Fig. 1.